



CONTRALLA

GROUP 27 FUNGICIDE

A fungicide for the control of potato blight (*Phytophthora Infestans*).

A water dispersible granule (WG) formulation containing 45% w/w cymoxanil.

SAFETY PRECAUTIONS

Operator Protection

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALL) AND SUITABLE PROTECTIVE GLOVES when handling the product.

WASH SPLASHES from skin immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking or smoking and after work.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

AVOID CONTACT WITH SKIN.

WEAR SUITABLE GLOVES.

Environmental Protection

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

Storage and Disposal

KEEP IN ORIGINAL CONTAINER tightly closed, in a safe place. EMPTY CONTAINER COMPLETELY and dispose of safely. KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

THIS MATERIAL AND ITS CONTAINER must be disposed of in a safe way.

PROTECT FROM FROST

UFI: A1Y2-W0M2-600J-54YS

SAFETY INFORMATION

FOR PROFESSIONAL USE ONLY

Warning



May cause an allergic skin reaction.

Causes serious eye irritation.

Suspected of damaging fertility or the unborn child.

May cause damage to organs through prolonged or repeated exposure (blood, thymus).

Very toxic to aquatic life with long lasting effects.

Do not breathe spray.

Avoid release to the environment.

Wear protective gloves.

If exposed or concerned: Get medical advice/attention.

Collect spillage.

Dispose of contents/container to a licensed hazardous waste disposal contractor except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS 06448

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crops/situations	Maximum individual dose (kg product/ha)	Maximum total dose per year (kg/ha)	Latest time of application
Potato	0.25	2.5	7 days before harvest

Control of late blight of potatoes caused by *phytophthora infestans* is achieved when used in conjunction with products containing active substances with multi-site activity.

Other specific restrictions:

The minimum interval between applications is 7 days.

Do not re-use container for any purpose.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

Net contents
4 x 5 kg e

Authorisation Holder and Marketing Company
JT Agro Europe sp. z o.o.
Gate A, Aleja Grunwaldza 472, 80-309 Gdańsk, Poland
Telephone: +44 1484 848885
www.jtcrop.com
info@jtcrop.com

